This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.



Bisphosphonate solution. Bone tumors induce osteoclast (tumor giant cells) formation Figure 1. A schematic diagram for therapeutic rationale using bone grafts coated with Osteoclasts destroy bone. Bisphosphonate-Bone Graft composite (🖏) can induce apoptosis in giant cell tumors (GCT) & other tumor cells, prevent recurrence and protect bone graft resorption.




Figure 2. Pamidronate induces apoptosis of tumor cells of giant cells in a dose dependent manner. The control picture shows plumpy, polyhedral cytoplasm. Addition of 100 & 200 uM of Pamidronate induces cell death. Annexin V staining indicates apoptosis of tumor cells.

